

Analytic solutions of relativistic hydrodynamics for a lattice QCD inspired equation of state

Máté Csanád, Márton Nagy, Sándor Lökös

Zimányi Winter School
Budapest, December 4, 2012

- 1 Hydrodynamics in relativistic heavy ion collisions
 - Basic equations and Equation of State
 - An already known solution
 - Observables with $\kappa = \text{const.}$

- 1 Hydrodynamics in relativistic heavy ion collisions
 - Basic equations and Equation of State
 - An already known solution
 - Observables with $\kappa = \text{const.}$
- 2 New solution for more general EoS
 - Temperature dependent EoS with conserved charge
 - Temperature dependent EoS without conserved charge
 - Pressure dependent EoS without conserved charge

- 1 Hydrodynamics in relativistic heavy ion collisions
 - Basic equations and Equation of State
 - An already known solution
 - Observables with $\kappa = \text{const.}$
- 2 New solution for more general EoS
 - Temperature dependent EoS with conserved charge
 - Temperature dependent EoS without conserved charge
 - Pressure dependent EoS without conserved charge
- 3 Investigation with lattice QCD parametrization
 - Temperature dependence from the new solution
 - Connection between initial and final state

Basic equations

- Continuity equation and energy-momentum equation:

$$\partial_\nu(nu^\nu) = 0 \quad , \quad \partial_\nu T^{\mu\nu} = 0.$$

- In a perfect fluid, $T^{\mu\nu} = (\epsilon + p)u^\mu u^\nu - g^{\mu\nu}p$.
- From the condition $\partial_\nu T^{\mu\nu} = 0$, the Euler equation and the energy conservation equation can be deduced:

$$\begin{aligned}(\epsilon + p)\partial_\nu u^\nu + u^\nu \partial_\nu \epsilon &= 0, \\ (\epsilon + p)u^\nu \partial_\nu u^\mu &= (g^{\mu\nu} - u^\mu u^\nu)\partial_\nu p.\end{aligned}$$

- EoS is $\epsilon = \kappa p$, where κ is not necessarily constant.

A known solution

(Csörgő, Csernai, Hama, Kodama, Heavy Ion Phys. A21:73-84, 2004)

- Ellipsoidal symmetry in space-time: $s = \frac{x^2}{X^2(t)} + \frac{y^2}{Y^2(t)} + \frac{z^2}{Z^2(t)}$

A known solution

(Csörgő, Csernai, Hama, Kodama, Heavy Ion Phys. A21:73-84, 2004)

- Ellipsoidal symmetry in space-time: $s = \frac{x^2}{X^2(t)} + \frac{y^2}{Y^2(t)} + \frac{z^2}{Z^2(t)}$
- Hubble-type velocity field: $u^\mu = \frac{x^\mu}{\tau} \quad \tau = \sqrt{x_\mu x^\mu}$

A known solution

(Csörgő, Csernai, Hama, Kodama, Heavy Ion Phys. A21:73-84, 2004)

- Ellipsoidal symmetry in space-time: $s = \frac{x^2}{X^2(t)} + \frac{y^2}{Y^2(t)} + \frac{z^2}{Z^2(t)}$
- Hubble-type velocity field: $u^\mu = \frac{x^\mu}{\tau} \quad \tau = \sqrt{X_\mu X^\mu}$
- The solution:
 - $n = n_0 \left(\frac{\tau_0}{\tau}\right)^3 \nu(s)$
 - $T = T_0 \left(\frac{\tau_0}{\tau}\right)^{3/\kappa} \frac{1}{\nu(s)}$
 - $p = nT$
- $\nu(s)$: arbitrary function of the s scale parameter (e.g. $e^{-bs/2}$)

A known solution

(Csörgő, Csernai, Hama, Kodama, Heavy Ion Phys. A21:73-84, 2004)

- Ellipsoidal symmetry in space-time: $s = \frac{x^2}{X^2(t)} + \frac{y^2}{Y^2(t)} + \frac{z^2}{Z^2(t)}$
- Hubble-type velocity field: $u^\mu = \frac{x^\mu}{\tau} \quad \tau = \sqrt{X_\mu X^\mu}$
- The solution:
 - $n = n_0 \left(\frac{\tau_0}{\tau}\right)^3 \nu(s)$
 - $T = T_0 \left(\frac{\tau_0}{\tau}\right)^{3/\kappa} \frac{1}{\nu(s)}$
 - $p = nT$
- $\nu(s)$: arbitrary function of the s scale parameter (e.g. $e^{-bs/2}$)
- Solution is a non-accelerating one

A known solution

(Csörgő, Csernai, Hama, Kodama, Heavy Ion Phys. A21:73-84, 2004)

- Ellipsoidal symmetry in space-time: $s = \frac{x^2}{X^2(t)} + \frac{y^2}{Y^2(t)} + \frac{z^2}{Z^2(t)}$
- Hubble-type velocity field: $u^\mu = \frac{x^\mu}{\tau} \quad \tau = \sqrt{X_\mu X^\mu}$
- The solution:
 - $n = n_0 \left(\frac{\tau_0}{\tau}\right)^3 \nu(s)$
 - $T = T_0 \left(\frac{\tau_0}{\tau}\right)^{3/\kappa} \frac{1}{\nu(s)}$
 - $p = nT$
- $\nu(s)$: arbitrary function of the s scale parameter (e.g. $e^{-bs/2}$)
- Solution is a non-accelerating one
- In this solution κ is constant

A known solution

(Csörgő, Csernai, Hama, Kodama, Heavy Ion Phys. A21:73-84, 2004)


- Ellipsoidal symmetry in space-time: $s = \frac{x^2}{X^2(t)} + \frac{y^2}{Y^2(t)} + \frac{z^2}{Z^2(t)}$
- Hubble-type velocity field: $u^\mu = \frac{x^\mu}{\tau} \quad \tau = \sqrt{X_\mu X^\mu}$
- The solution:
 - $n = n_0 \left(\frac{\tau_0}{\tau}\right)^3 \nu(s)$
 - $T = T_0 \left(\frac{\tau_0}{\tau}\right)^{3/\kappa} \frac{1}{\nu(s)}$
 - $p = nT$
- $\nu(s)$: arbitrary function of the s scale parameter (e.g. $e^{-bs/2}$)
- Solution is a non-accelerating one
- In this solution κ is constant
- Entropy density σ can be calculated as well

Observables with $\kappa = \text{constant}$

- Source function can be calculated from this solution


$$S(x, p) d^4x = \mathcal{N} n(x) p^\mu d^3\Sigma_\mu(x) H(\tau) d\tau \exp\left(-\frac{p_\mu u^\mu}{T}\right)$$

- Calculated observables are fitted to data successfully
Csanád, Vargyas, Eur. Phys. J. A 44, 473 (2010), arXiv:0909.4842




More general EoS

- Constant EoS may not be realistic (κ may depend on temperature or pressure)




More general EoS

- Constant EoS may not be realistic (κ may depend on temperature or pressure)
- From IQCD trace anomaly ($I = \epsilon - 3p$) can be calculated
Pressure is given by $\frac{p(T)}{T^4} = \int_0^T \frac{dT}{T} \frac{I(T)}{T^4}$



More general EoS

- Constant EoS may not be realistic (κ may depend on temperature or pressure)
- From IQCD trace anomaly ($I = \epsilon - 3p$) can be calculated
Pressure is given by $\frac{p(T)}{T^4} = \int_0^T \frac{dT}{T} \frac{I(T)}{T^4}$
- From $I \rightarrow \kappa(T) = I(T)/p(T) + 3$, speed of sound: $\kappa = \frac{1}{c_s^2}$



Temperature dependent EoS with conserved charge

- If there is a conserved charge (n), the energy equation yields


$$T \partial_\nu u^\nu + \frac{d}{dT} (T \kappa(T)) u^\nu \partial_\nu T = 0.$$

Temperature dependent EoS with conserved charge

- If there is a conserved charge (n), the energy equation yields

$$T \partial_\nu u^\nu + \frac{d}{dT} (T \kappa(T)) u^\nu \partial_\nu T = 0.$$

- It works only if $d(T\kappa(T))/dT \geq 0$




Temperature dependent EoS with conserved charge

- If there is a conserved charge (n), the energy equation yields

$$T \partial_\nu u^\nu + \frac{d}{dT} (T \kappa(T)) u^\nu \partial_\nu T = 0.$$

- It works only if $d(T \kappa(T)) / dT \geq 0$



- Solution cannot be applied if $173 \text{ MeV} < T < 225 \text{ MeV}$
- This problem is absent, if there is no conserved charge

Temperature dependent EoS without conserved charge

- If there is no conserved charge, we have

$$\epsilon = Ts - p \rightarrow d\epsilon = Tds. \quad (1)$$

(because of Gibbs–Duhem relation: $dp = sdT$)

Temperature dependent EoS without conserved charge

- If there is no conserved charge, we have

$$\epsilon = Ts - p \rightarrow d\epsilon = Tds. \quad (1)$$

(because of Gibbs–Duhem relation: $dp = sdT$)

- Put (1) and $\epsilon + p = Ts$ to the energy equation:

$$T\sigma\partial_\nu u^\nu + u^\nu T\partial_\nu\sigma \rightarrow \partial_\nu(\sigma u^\nu) = 0$$

This is a continuity equation to entropy-density.

Temperature dependent EoS without conserved charge

- If there is no conserved charge, we have

$$\epsilon = Ts - p \rightarrow d\epsilon = Tds. \quad (1)$$

(because of Gibbs–Duhem relation: $dp = sdT$)

- Put (1) and $\epsilon + p = Ts$ to the energy equation:

$$T\sigma\partial_\nu u^\nu + u^\nu T\partial_\nu\sigma \rightarrow \partial_\nu(\sigma u^\nu) = 0$$

This is a continuity equation to entropy-density.

- Put $\epsilon + p = Ts$ and $\epsilon = \kappa(T)p$ to the energy equation

$$T\partial_\nu u^\nu + \left(\kappa + \frac{T}{\kappa + 1} \frac{d\kappa(T)}{dT} \right) u^\nu \partial_\nu T = 0$$

Temperature dependent EoS without conserved charge

- If there is no conserved charge, we have

$$\epsilon = Ts - p \rightarrow d\epsilon = Tds. \quad (1)$$

(because of Gibbs–Duhem relation: $dp = sdT$)

- Put (1) and $\epsilon + p = Ts$ to the energy equation:

$$T\sigma\partial_\nu u^\nu + u^\nu T\partial_\nu\sigma \rightarrow \partial_\nu(\sigma u^\nu) = 0$$

This is a continuity equation to entropy-density.

- Put $\epsilon + p = Ts$ and $\epsilon = \kappa(T)p$ to the energy equation

$$T\partial_\nu u^\nu + \left(\kappa + \frac{T}{\kappa + 1} \frac{d\kappa(T)}{dT} \right) u^\nu \partial_\nu T = 0$$

- It's not the same equation (only if $\kappa = \text{const.}$)

New solution with more general EoS

Csanád, Nagy, Lökös, Eur. Phys. J. A (2012) 48: 173

If there is a conserved charge

$$n = n_0 \frac{\tau_0^3}{\tau^3},$$
$$u^\nu = \frac{x^\nu}{\tau},$$
$$\frac{\tau_0^3}{\tau^3} = \exp \left[\int_{T_0}^T \frac{d\kappa(\zeta)\zeta}{d\zeta} \frac{1}{\zeta} d\zeta \right]$$

New solution with more general EoS

Csanád, Nagy, Lökös, Eur. Phys. J. A (2012) 48: 173

If there is a conserved charge

$$n = n_0 \frac{\tau_0^3}{\tau^3},$$
$$u^\nu = \frac{x^\nu}{\tau},$$
$$\frac{\tau_0^3}{\tau^3} = \exp \left[\int_{T_0}^T \frac{d\kappa(\zeta)\zeta}{d\zeta} \frac{1}{\zeta} d\zeta \right]$$

- Arbitrary $\kappa(T)$ function may be used
- If $d\kappa(T)T/dT < 0 \rightarrow \partial_\nu u^\nu < 0!$ It's not realistic!

New solution with more general EoS

Csanád, Nagy, Lökös, Eur. Phys. J. A (2012) 48: 173

If there is no conserved charge

$$\begin{aligned}\sigma &= \sigma_0 \frac{\tau_0^3}{\tau^3}, \\ u^\nu &= \frac{x^\nu}{\tau}, \\ \frac{\tau_0^3}{\tau^3} &= \exp \left[\int_{T_0}^T \left(\frac{\kappa(\zeta)}{\zeta} + \frac{1}{\kappa + 1} \frac{d\kappa(\zeta)}{d\zeta} d\zeta \right) \right]\end{aligned}$$

- It can be used if κ is given as a function of temperature

Pressure dependent κ without conserved charge

Csanád, Nagy, Lökös, Eur. Phys. J. A (2012) 48: 173

- If EoS is given as a function of pressure: $\epsilon = \kappa(p)p$

Pressure dependent κ without conserved charge

Csanád, Nagy, Lökös, Eur. Phys. J. A (2012) 48: 173

- If EoS is given as a function of pressure: $\epsilon = \kappa(p)p$
- The energy equation can be written with this EoS as

$$\partial_\nu(\epsilon u^\nu) + p\partial_\nu u^\nu = 0$$

Pressure dependent κ without conserved charge

Csanád, Nagy, Lökös, Eur. Phys. J. A (2012) 48: 173

- If EoS is given as a function of pressure: $\epsilon = \kappa(p)p$
- The energy equation can be written with this EoS as

$$\partial_\nu(\epsilon u^\nu) + p\partial_\nu u^\nu = 0$$

- From this equation

$$\left(\frac{\epsilon}{p} + 1\right) \partial_\nu \ln\left(\frac{V_0}{V}\right) = \frac{\partial_\nu \epsilon}{p}$$

Pressure dependent κ without conserved charge

Csanád, Nagy, Lökös, Eur. Phys. J. A (2012) 48: 173

- If EoS is given as a function of pressure: $\epsilon = \kappa(p)\rho$
- The energy equation can be written with this EoS as

$$\partial_\nu(\epsilon u^\nu) + p\partial_\nu u^\nu = 0$$

- From this equation

$$\left(\frac{\epsilon}{p} + 1\right) \partial_\nu \ln\left(\frac{V_0}{V}\right) = \frac{\partial_\nu \epsilon}{p}$$

- With an integral transformation it can be „solved” by

$$\frac{V_0}{V} = \exp\left[\int_{p_0}^p \left(\frac{\kappa(\zeta)}{\zeta} + \frac{d\kappa}{d\zeta}\right) \frac{d\zeta}{\kappa(\zeta) + 1}\right]$$

A new solution if $\kappa = \kappa(p)$

Csanád, Nagy, Lökös, Eur. Phys. J. A (2012) 48: 173

This new solution can be written as


$$\begin{aligned}\sigma &= \sigma_0 \frac{\tau_0^3}{\tau^3}, \\ u^\nu &= \frac{x^\nu}{\tau}, \\ \frac{\tau_0^3}{\tau^3} &= \exp \left[\int_{p_0}^p \left(\frac{\kappa(\zeta)}{\zeta} + \frac{d\kappa}{d\zeta} \right) \frac{d\zeta}{\kappa(\zeta) + 1} \right]\end{aligned}$$

- It can be used if a parametrization to $\epsilon(p)$ is given
- If $p = p(T)$, we get the previous solution

Investigation without conserved charge

- Let assume IQCD EoS
- $T(\tau)$ can be calculated

$$\frac{V_0}{V} = \exp \left[\int_{T_0}^T \left(\frac{\kappa(\zeta)}{\zeta} + \frac{1}{\kappa + 1} \frac{d\kappa(\zeta)}{d\zeta} d\zeta \right) \right]$$



Summary

- Constant EoS is succesful in description of data

Summary

- Constant EoS is succesful in description of data
- According to IQCD calculation, EoS is not constant

Summary

- Constant EoS is succesful in description of data
- According to IQCD calculation, EoS is not constant
- Found new solutions with temperature, pressure dependent EoS

Summary

- Constant EoS is succesful in description of data
- According to IQCD calculation, EoS is not constant
- Found new solutions with temperature, pressure dependent EoS
- $T(\tau)$ can be calculated \rightarrow IQCD EoS applicable

Summary

- Constant EoS is succesful in description of data
- According to IQCD calculation, EoS is not constant
- Found new solutions with temperature, pressure dependent EoS
- $T(\tau)$ can be calculated \rightarrow IQCD EoS applicable
- If assuming τ_f/τ_{init} , T_f/T_{init} can be calculated

Thank you for your attention!